




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 413/10, A61K 31/42, C07D 413/14	A1	(11) International Publication Number: WO 99/24428 (43) International Publication Date: 20 May 1999 (20.05.99)
(21) International Application Number: PCT/US98/22639 (22) International Filing Date: 30 October 1998 (30.10.98) (30) Priority Data: 60/065,376 12 November 1997 (12.11.97) US (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): HESTER, Jackson, B., Jr. [US/US]; 9219 East ML Avenue, Galesburg, MI 49053 (US). (74) Agent: PHARMACIA & UPJOHN COMPANY; Intellectual Property Legal Services, 301 Henrietta Street, Kalamazoo, MI 49001 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: OXAZOLIDINONE DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS



(I)

(57) Abstract

A compound of Formula (I) or a pharmaceutically acceptable salt thereof, which is antimicrobial agents, effective against various human and veterinary pathogens, including gram positive aerobic organisms, gram negative organisms, and anaerobic organisms.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

OXAZOLIDINONE DERIVATIVES AND PHARMACEUTICAL COMPOSITIONS

BACKGROUND OF THE INVENTION

The present invention relates to novel oxazolidinone compounds or
5 pharmaceutically acceptable salts thereof, and pharmaceutical agents that contain them as
active ingredients for preventing or treating infectious diseases. The compounds are
unique oxazolidinones having a hexahydro-1,4-diazepin-5-one substituent.

More specifically, novel oxazolidinone compounds of the present invention
relates to useful antimicrobial agents, effective against various human and veterinary
10 pathogens, including gram positive aerobic organisms such as multiply-resistant
staphylococci and *streptococci*, gram negative organisms such as *H. influenzae* and *M.*
catarrhalis as well as anaerobic organisms such as *bacteroides* and *clostridia* species,
and acid-resistant organisms such as *Mycobacterium tuberculosis* and *Mycobacterium*
avium.

15

INFORMATION DISCLOSURE

International Publication No. 97/27188 discloses piperazine-3-one analogs which
are homologs of the invention.

International Publication No. WO93/23384 discloses oxazolidinones containing a
20 substituted diazine (piperazine) moiety and their uses as antimicrobials.

International Publication No. WO93/09103 discloses substituted aryl and
heteroaryl-phenyl-oxazolidinones useful as antimicrobials.

International Publication No. WO90/02744 discloses 5'-indoliny-5-
amidomethyloxazolidinones, 3-(fused-ring substituted)phenyl-5-
25 midomethyloxazolidinones, and 3-(nitrogen substituted)-phenyl-5-
amidomethyloxazolidinones which are useful as antibacterial agents.

Other references disclosing various oxazolidinones include US Patents 5,547,950,
4,801,600, J. Med. Chem., 32, 1673-81 (1989); J. Med. Chem., 33, 2569-78 (1990);
Tetrahedron, 45, 1323-26 (1989); and J. Med. Chem., 35, 1156 (1992).


30 European Patent Publication 352,781 discloses phenyl and pyridyl substituted
phenyl oxazolidinones.

European Patent Publication 316,594 discloses 3-substituted styryl
oxazolidinones.

European Patent Publication 312,000 discloses phenylmethyl and pyridylmethyl substituted phenyl oxazolidinones.

SUMMARY OF THE INVENTION

The present invention provides an oxazolidinone derivative represented by the general structural Formula I or a pharmaceutically acceptable salt thereof



I

or a pharmaceutically acceptable salt thereof wherein:

R is H, C₂₋₆ alkenyl, C₂₋₇ alkynyl, C₁₋₆ alkyl, or C₁₋₆ alkyl substituted with one or two of the following:

- a) F,
- b) Cl,
- c) CF₃,
- d) -OH,
- e) C₁₋₄ alkoxy,
- f) -CH₂C(=O)C₁₋₄ alkyl,
- g) -OC(=O)N(R₄)₂,
- h) C₁₋₄ alkyl S(O)_n, (wherein n is 0 to 2),
- i) -CN,
- j) carboxy,
- k) -C₁₋₄ alkoxy carbonyl,
- l) -C(=O)N(R₄)₂,
- m) -N(R₄)SO₂C₁₋₄ alkyl,
- n) -N(R₄)C(=O)C₁₋₄ alkyl,
- o) -N(R₄)C(=O)N(R₄)₂,
- p) -N(R₄)C(=O)C₁₋₄ alkoxy,
- q) aryl, or

r) Het;

aryl is phenyl, optionally substituted with one or two of the following:

- a) F,
- b) Cl,
- 5 c) Br,
- d) -CF₃,
- e) CN,
- f) C₁₋₃ alkoxy, or
- g) C₁₋₃ alkylthio;


10 Het is a 5- or 6-membered heteroaromatic moiety having 1-3 N, O or S atoms, optionally substituted with the following:

- a) F,
- b) Cl,
- c) C₁₋₃ alkoxy,
- 15 d) C₁₋₃ alkylthio, or
- e) CN;

R₁ and R₂ are independently H, F, or Cl;

R₃ is

- a) C₁₋₆ alkyl, optionally substituted with one to three F or one to two Cl,
- 20 b) C₁₋₆ alkoxy,
- c) amino,
- d) C₁₋₆ alkylamino,
- e) C₁₋₆ dialkylamino
- f) C₃₋₆ cycloalkyl,
- 25 g) C₁₋₆ alkylthio, or

h)  (wherein m is 0, 1, 2, 3 or 4);

R₄ is

- 30 a) H, or
- b) C₁₋₃ alkyl; and

X is O or S.

The present invention also provides an antimicrobial agent or pharmaceutical composition that contains the oxazolidinone compound or a pharmaceutically acceptable salt thereof as an active ingredient. The antimicrobial agent containing the active ingredient of the present invention can be used for treatment or prevention of infectious diseases.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of Formula I as structurally disclosed above are useful antimicrobials. Typically, as further explained below, the compounds can be administered as antibacterial agents in a dosage range of from about 0.1 to 100 mg/kg or preferably from about 3.0 to about 50 mg/kg of body weight per day.

In the structural formula shown above the carbon content of various hydrocarbon containing moieties is indicated by a prefix designating the minimum and maximum number of carbon atoms in the moiety, i.e., the prefix C_i-C_j defines the number of carbon atoms present from the integer "i" to the integer "j" inclusive.

The term "C₁₋₆ alkyl" used herein refers to an alkyl group having one to six carbon atoms such as, for example, methyl, ethyl, propyl, butyl, pentyl, hexyl and isomeric forms thereof; preferably methyl, ethyl, propyl and isomeric forms thereof.

The term "C₂₋₆ alkenyl" refers to at least one double bound alkenyl group having two to six carbon atoms such as, for example, vinyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 1-pentenyl, 2-pentenyl, 1-hexenyl and isomeric forms thereof, preferably an alkenyl group having 2 to 6 carbon atoms, and more preferably an alkenyl group having 2 to 4 carbon atoms.

The term "C₂₋₇ alkynyl" refers to at least one triple bond alkynyl group having two to seven carbon atoms such as, for example, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl and their isomeric forms thereof.

The term "C₁₋₆ alkylamino" refers to an alkyl group having one to six carbon atoms attached to an amino moiety.

The term "C₁₋₆ dialkylamino" refers to two alkyl groups having one to six carbon atoms attached to an amino moiety.

The term "C₁₋₄ alkoxy" refers to an alkyl group having one to four carbon atoms attached an oxygen atom of hydroxyl group such as, for example, methoxy, ethoxy,

propoxy, butoxy and isomeric forms thereof, preferably an alkoxy group having 1 to 2 carbon atoms.

The term "C₁₋₆ alkylthio" refers to an alkyl group having one to six carbon atoms attached a thio moiety such as, for example, methylthio, ethylthio, propylthio and isomeric forms thereof, preferably an alkylthio group having 1 to 2 carbon atoms.

The term "C₃₋₆ cycloalkyl" refers to three to six carbon atoms forming cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and isomeric forms thereof.

The term "aryl" refers to an phenyl moiety optionally substituted with one or two F, Cl, Br, -CF₃, -CN, -C₁₋₃ alkoxy, or -C₁₋₃ alkylthio;

The term "Het" refers to a 5- or 6-membered heteroaromatic moiety having one to three atoms selected from the group consisting of O, N or S atoms such as, for example, furan, thiophene, pyrrole, pyrazole, triazoles, oxazole, thiazole, isothiazole, oxadiazoles, oxathiazole, pyridine, pyridazine, pyrimidine, pyrazine, piperazine and triazines all of which can optionally be substituted with one substituent selected from the group consisting of F, Cl, C₁₋₃ alkoxy, C₁₋₃ alkylthio or CN.

The compounds of the present invention can be converted to their salts according to conventional methods.

Pharmaceutically acceptable salts means acid addition salts useful for administering the compounds of this invention and these include hydrochloride, hydrobromide, sulfate, phosphate, acetate, propionate, lactate, mesylate, maleate, succinate, tartrate, citrate, 2-hydroxyethyl sulfonate, fumarate and the like when a basic group is present. These salts may be in hydrated form. Some of the compounds of this invention may form metal salts such as sodium, potassium, calcium and magnesium salts and these are embraced by the term "pharmaceutically acceptable salts".


Due to the configuration at C-5 of the oxazolidinone ring of compounds as represented in the structure of Formula I the compounds of this invention may exist in geometric, optical and other isomeric forms and this invention embraces any of these isomers. The racemic mixture and enantiomers are all believed to be useful as an antibacterial. Regardless, the preferred absolute configuration at C-5 of the oxazolidinone ring of compounds is as represented in the structure of Formula I. This absolute configuration is called (S) under the Cahn-Ingold-Prelog nomenclature system. It is believed that a majority of the pharmacological activity resides in this (S)-enantiomer to produce the antibacterial effect.

Compounds of Formula I can be prepared as shown in Scheme I where P represents an alcohol protecting group such as benzyl or *tert*-butyldimethylsilyl. Structure 2 of this scheme are prepared according to the methods outlined in Example 1, Step 1 and 2. In Scheme I, the alcohols of 2 are protected as benzyl ethers. In a suitable procedure for this reaction, a solution of the alcohol 2 in a solvent such as Et₂O or THF is allowed to react first with sodium hydride at 0-25 °C and then with benzyl bromide and tetrabutylammonium iodide at 0-25 °C to give structure 3. The ethylene ketal of 3 can then be removed with an acidic catalyst such as *p*-toluenesulfonic acid in acetone (as described in Example 1, Step 2) to give structure 5 where P is benzyl. Alternatively the ketal of 2 can be removed, the resulting structure 4 is allowed to react with *tert*-butyldimethylsilyl chloride and imidazole in DMF or *tert*-butyldimethylsilyl chloride and triethyl amine in methylene chloride to give structure 5 where the alcohol protecting group (P) is *tert*-butyldimethylsilyl. Ketone 5 is then allowed to react with hydroxylamine hydrochloride and sodium acetate in methanol-methylene chloride to give oxime 6 (see Example 1, Step 3). The Beckmann rearrangement of structure 6 is carried out with *p*-toluenesulfonyl chloride and sodium carbonate in aqueous acetone at 20-40 °C to give structure 7. For compounds of Formula I where R is not hydrogen, compounds 7 can be alkylated with R'Y where Y is Br, I, CH₃SO₃ or *p*-CH₃PhSO₃ and R' is an appropriate alkyl substituent. In one method for this alkylation compounds of structure 7 are allowed to react with sodium hydride and R'Y in a solvent such as DMF at 0-25 °C to give 8. Alternatively, structure 7 can react with R'Y, potassium hydroxide and tetrabutylammonium bromide in THF or acetonitrile at 20-50 °C to give 8. Deprotection of the alcohols 7 or 8 provide structure 9. When P is a benzyl ether, this can be accomplished by hydrogenolysis with hydrogen and a palladium catalyst in ethanol or with ammonium formate and a palladium catalyst in methanol at 10-30 °C. The *tert*-butyldimethylsilyl protecting group can be removed under acidic conditions or with fluoride ion. This deprotection can be carried out, for example, with trifluoroacetic acid in methylene chloride at 25 °C or with tetrabutylammonium fluoride in THF at 25 °C to give alcohol 9. Transformation of the alcohol 9 to the amine 11 can be carried out as described in Example 1, Step 1. Alternatively, the reaction of 9 with *m*-nitrobenzenesulfonyl chloride and triethylamine in methylene chloride at 5-25 °C will give the *m*-nitrobenzenesulfonate 10 which will react with ammonium hydroxide in THF or acetonitrile -isopropanol at 30-60 °C to give the amine 11. The reaction of compound

11 with acyl halides, anhydrides, isocyanates, isothiocyanates or dithioesters provides compounds of Formula I.

Compounds of Formula I where R is hydrogen and X is oxygen are conveniently prepared by allowing compounds **12** to react with p-toluenesulfonyl chloride and sodium carbonate in aqueous acetone at 20-40 °C (see Example 1, Step 4).

Scheme I



The compounds of the invention are useful for the treatment of microbial infections in humans and other warm blooded animals by either parenteral, oral, or topical administration.

The term "treatment" as used herein means partial or total lessening of symptoms of a disease which a patient suffers from; the term "prevention" as used herein means partial or total avoidance of symptoms of a disease in a patient who, according to a doctor's diagnosis, may suffer from the disease or a related state unless the preventive measure is taken.

The pharmaceutical compositions of this invention may be prepared by combining the compounds of Formula I of this invention with a solid or liquid pharmaceutically acceptable carrier and, optionally, with pharmaceutically acceptable adjuvants and excipients employing standard and conventional techniques. Solid form compositions include powders, tablets, dispersible granules, capsules and suppositories. A solid carrier can be at least one substance which may also function as a diluent, flavoring agent, solubilizer, lubricant, suspending agent, binder, tablet disintegrating agent, and encapsulating agent. Inert solid carriers include magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, cellulosic materials, low melting wax, cocoa butter, and the like. Liquid form compositions include solutions, suspensions and emulsions. For example, there may be provided solutions of the compounds of this invention dissolved in water and water-propylene glycol and water-polyethylene glycol systems, optionally containing conventional coloring agents, flavoring agents, stabilizers and thickening agents.

Preferably, the pharmaceutical composition is made by employing conventional techniques in unit dosage form containing effective amounts of the active component, that is, the compound of Formula I according to this invention.

The quantity of active component, that is the compound of Formula I, in the pharmaceutical composition and unit dosage form thereof may be varied or adjusted widely depending upon the particular application method, the potency of the particular compound and the desired concentration. Generally, the quantity of active component will range between 0.5% to 90% by weight of the composition.

In therapeutic use for treating, or combatting bacterial infections in humans and other animals that have been diagnosed with bacterial infections, the compounds or pharmaceutical compositions thereof will be administered orally, parenterally and/or

topically at a dosage to obtain and maintain a concentration, that is, an amount, or blood-level of active component in the animal undergoing treatment which will be antibacterially effective. Generally, such antibacterially effective amount of dosage of active component will be in the range of about 0.1 to about 100 mg/kg, more preferably about 3.0 to about 50 mg/kg of body weight/day. It is to be understood that the dosages may vary depending upon the requirements of the patient, the severity of the bacterial infection being treated, and the particular compound being used. Also, it is to be understood that the initial dosage administered may be increased beyond the above upper level in order to rapidly achieve the desired blood-level or the initial dosage may be smaller than the optimum and the daily dosage may be progressively increased during the course of treatment depending on the particular situation. If desired, the daily dose may also be divided into multiple doses for administration, e.g., two to four times per day.

The compounds of Formula I are administered parenterally, i.e., by injection, for example, by intravenous injection or by other parenteral routes of administration.

Pharmaceutical compositions for parenteral administration will generally contain a pharmaceutically acceptable amount of the compound according to Formula I as a soluble salt (acid addition salt or base salt) dissolved in a pharmaceutically acceptable liquid carrier such as, for example, water-for-injection and a suitably buffered isotonic solution, for example, having a pH of about 3.5-6. Suitable buffering agents include, for example, trisodium orthophosphate, sodium bicarbonate, sodium citrate, N-methylglucamine, L(+)-lysine and L(+)-arginine, to name a few. The compound according to Formula I generally will be dissolved in the carrier in an amount sufficient to provide a pharmaceutically acceptable injectable concentration in the range of about 1 mg/ml to about 400 mg/ml. The resulting liquid pharmaceutical composition will be administered so as to obtain the above-mentioned antibacterially effective amount of dosage. The compounds of Formula I according to this invention are advantageously administered orally in solid and liquid dosage forms.


As a topical treatment, an effective amount of a compound of Formula I is admixed in a pharmaceutically acceptable gel or cream vehicle that can be applied to the patient's skin at the area of treatment. Preparation of such creams and gels is well known in the art and can include penetration enhancers.

The compounds of this invention are useful antimicrobial agents, effective against various human and veterinary pathogens, including gram positive aerobic organisms such

as multiply-resistant *staphylococci* and *streptococci*, gram negative organisms such as *H. influenzae* and *M. catarrhalis* as well as anaerobic organisms such as *bacteroides* and *clostridia* species, and acid-resistant organisms such as *Mycobacterium tuberculosis* and *Mycobacterium avium*.

In order to more fully illustrate the nature of the invention and the manner of practice the same, the following experimental examples are presented, but they should not be taken as limitations.

EXAMPLE 1 Preparation of (S)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide.



Step 1: Preparation of (S)-N-[3-(3-fluoro-4-piperidin-1-yl-phenyl)-2-oxo-oxazolidin-5-ylmethyl]-acetamide :

Diisopropylethylamine (15.7 ml) and 3,4-difluoronitrobenzene (5.0 ml) are added successively to an ethyl acetate solution (70 ml) of piperidine (5.77 g) and the mixture is stirred at room temperature for 2 days. Water is added to the reaction solution and the separating ethyl acetate layer are washed with water and brine, dried over anhydrous sodium sulfate. The solvent is evaporated to afford a nitro compound (10.1 g) in a yield of 100%. Palladium on carbon (10%, 1.0 g) is added to an ethyl acetate solution (101 ml) of the nitro compound (10.1 g) and the mixture is stirred at room temperature for 4 hours under hydrogen atmosphere. The palladium on carbon is filtered off and the filtrate is concentrated under vacuum to yield an amine (8.75 g, 100%). Sodium hydrogencarbonate (5.0 g) and benzyloxycarbonyl chloride (8.4 ml) are added successively to a tetrahydrofuran (THF) solution (100 ml) of the amine (8.75 g), and the mixture is stirred at room temperature for 14 hours. Water is added to the reaction solution and the separating THF layer is washed with water and brine, dried over anhydrous sodium sulfate. The solvent is evaporated and the residue is purified by silica gel column chromatography (solvent: ethyl acetate/hexane/chloroform = 1/6/4) to afford a benzyl carbamate (14.5 g) in a yield of 98%. Butyl lithium (1.6 M hexane solution: 5.2

ml) is added to a THF solution (24 ml) of the benzyl carbamate (2.75 g) at -78 C and the mixture is stirred for 5 min. At the same temperature, (R)-(-)-glycidyl butyrate (1.25 ml) is added to the stirred solution and the mixture is stirred for 14 hours while the temperature is raised slowly to room temperature. Water is added to the reaction solution and the separating THF layer is washed with water and brine, dried over anhydrous sodium sulfate. The solvent is evaporated and the residue is purified by silica gel column chromatography (solvent: ethyl acetate/hexane = 3/1) to afford an alcohol (2.20 g) in a yield of 89%. Tosyl chloride (2.85 g) is added to a pyridine solution (8 ml) of the alcohol (2.20 g) and the mixture is stirred at room temperature for 6 h. Water (32 ml) is added to the reaction solution and the mixture is stirred for 1 hour. The resulting precipitate is collected by filtration and washed with water, followed by drying under vacuum at room temperature to afford a tosylate (3.28 g) in a yield of 98%. Sodium azide (3.80 g) is added to a dimethylformamide (DMF) solution (23 ml) of the tosylate (3.28 g) at room temperature and the mixture is stirred at 65 C for 5.5 hours. After the reaction mixture is cooled to room temperature, water is added and the mixture is extracted with ethyl acetate; the organic layer is concentrated under vacuum. The resulting residue is dissolved in ethyl acetate and washed with water and brine, dried over anhydrous sodium sulfate. The solvent is evaporated and the residue is purified by silica gel column chromatography (solvent: ethyl acetate/hexane = 1/1) to afford an azide (2.20 g) in a yield of 94%. Acetic anhydride (0.65 ml) and pyridine (1.0 ml) are added to an ethyl acetate solution (19 ml) of the azide (2.20 g) at room temperature; after addition of palladium on carbon (10%, 0.22 g), the mixture is stirred at room temperature for 6 hours under 1 atm hydrogen atmosphere. The palladium on carbon is filtered off and the filtrate is washed with water and brine, dried over anhydrous sodium sulfate. The solvent is evaporated and the residue is purified by silica gel column chromatography (solvent: acetone/hexane = 1/1) to afford the title compound.

Step 2: Preparation of (S)-N-{3-[3-fluoro-4-(4-oxo-piperidin-1-yl)-phenyl]-2-oxo-oxazolidin-5-ylmethyl}-acetamide.

Using a commercially available 1,4-dioxo-8-aza-spiro[4.5]decane, (S)-N-{3-[4-(1,4-dioxo-8-aza-spiro[4.5]dec-8-yl)-3-fluoro-phenyl]-2-oxo-oxazolidin-5-ylmethyl}-acetamide is synthesized by the same method as in Step 1. To an acetone solution (70 ml) of this compound (3.79 g), water (20 ml) and p-toluenesulfonic acid monohydrate (3.66 g) are added successively and the mixture is heated under reflux for 3 hours. After

the reaction mixture is cooled to room temperature, acetone is distilled off and the aqueous layer is neutralized with triethylamine. The solution is extracted with methylene chloride and the organic layer is washed with brine, dried over anhydrous sodium sulfate. The solvent is evaporated and the residue is purified by silica gel column chromatography (solvent: chloroform/methanol = 50/1 - 25/1) to afford the title compound .

Step 3: Preparation of (S)-N-{3-[3-fluoro-4-(4-hydroxyimino-piperidin-1-yl)-phenyl]-2-oxo oxazolidin-5-ylmethyl}-acetamide.

Sodium acetate (517 mg) and hydroxylamine hydrochloride (219 mg) are successively added to a methanol-methylene chloride solution (10-10 ml) of 1.00 g of the product of Step 2, and the mixture is stirred at room temperature for 2 days. The solvent is evaporated and the residue is dissolved in methanol, followed by addition of a silica gel (8 g). Methanol is evaporated and the residue is purified by silica gel column chromatography (solvent: chloroform/methanol = 50/1 - 25/1) to afford the title compound.

Step 4: Preparation of (S)-N-[[3-[3-Fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide.


A stirred mixture of the compound of product of Step 3 (0.200 g, 0.549 mmol) in acetone (5.3 mL), under nitrogen, is treated first with 5% aqueous sodium carbonate (5.3 mL) and then, dropwise during 3 minutes with a solution of *p*-toluenesulfonyl chloride (0.16 g, 0.82 mmol) in acetone (2.7 mL). Initially this mixture is a two phase solution; however, after about 25 minutes a precipitate began to form. It is kept at ambient temperature (23 °C) for 4 hours and filtered. The filtrate is concentrated under reduced pressure to remove acetone and the aqueous residue is extracted with CH₂Cl₂. The extract is dried (MgSO₄) and concentrated to give a small amount of crude product. Most of the product is in the aqueous layer which is concentrated in vacuo. The residue is combined with the crude product from the CH₂Cl₂ extract and chromatographed on silica gel with mixtures of MeOH-NH₄OH-CH₂Cl₂ that continued 3-5% MeOH and 0.3-0.5% NH₄OH. The product is crystallized from MeOH-EtOAc to give the title compound. mp 140-146 °C;

MS *m/z* (relative intensity) 364 (M⁺, 96.1), 320 (100), 306 (6.7), 294 (10.9), 236 (41.8);

HRMS calcd for C₁₇H₂₁FN₄O₄; 364.1547 (M⁺); found 364.1545;

¹H NMR [300 MHz, (CD₃)₂SO] δ 1.81 (s, 3H), 2.57 (m, 2H), 3.07 (m, 4H), 3.24 (m, 2H), 3.38 (t, 2H), 3.67 (d, d, 1H), 4.06 (t, 1H), 4.68 (m, 1H), 7.08 (t, 1H), 7.13 (d, d, 1H), 7.45 (d, d, 1H), 7.65 (t, 1H), 8.21 (t, 1H).

EXAMPLE 2: Preparation of (S)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]thioacetamide.



Step 1: Preparation of (S)-[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)-phenyl]-2-oxo-5-oxazolidinyl]methyl *tert*-butyldimethylsilyl ether.

A stirred solution of 10.6 g (0.03 mol) of (S)-[3-[4-(1,4-dioxo-8-azaspiro[4.5]dec-8-yl)-3-fluorophenyl]-2-oxo-5-oxazolidinyl]methanol, the intermediate of formula 2 (Scheme 1) for the preparation of (S)-N-{3-[3-fluoro-4-(4-oxopiperidin-1-yl)phenyl]-2-oxooxazolidin-5-ylmethyl}-acetamide (Example 1, Step 2), in acetone (230 mL) is treated with water (65 mL) and *p*-toluenesulfonic acid monohydrate (11.4 g, 0.06 mol), refluxed under nitrogen for 5 hours and kept at ambient temperature (24 °C) for 18 hours. It is then concentrated in vacuo to remove acetone. The aqueous residue is neutralized with sodium bicarbonate and extracted with ethyl acetate; the extract is washed with saturated sodium bicarbonate, water and dilute sodium chloride, dried (Na₂SO₄) and concentrated to give the ketone, a compound of formula 4 (Scheme 1). A stirred solution of the ketone and triethylamine (12.5 mL, 0.09 mol) in methylene chloride (100 mL) is treated with *tert*-butyldimethylsilyl chloride (6.03 g, 0.04 mol) and kept under nitrogen at ambient temperature for 23 hours. Additional *tert*-butyldimethylsilyl chloride (3.0 g) is added and the mixture is kept at ambient temperature for an additional 20 hours. Additional triethylamine (3.0 mL) and *tert*-butyldimethylsilyl chloride (3.0 g) are again added and the mixture is kept at ambient temperature for 4 days, diluted with methylene chloride, washed with water and dilute sodium chloride, dried (Na₂SO₄) and concentrated. Chromatography of the residue on silica gel with mixtures of acetone-

heptane that contained 20-30% acetone gave 7.72 g of the *tert*-butyldimethylsilyl (TBDMS) ether, a compound of formula 5 (Scheme 1) where P is TBDMS. A stirred solution of the TBDMS ether (7.27 g, 17.2 mmol) in methanol (150 mL) is treated dropwise with a solution of hydroxylamine hydrochloride (1.44 g, 0.021 mol) and sodium acetate (1.72 g, 0.021 mol) in water (15 mL) and kept at ambient temperature for 20 hours. The mixture is concentrated under reduced pressure. A solution of the residue, a white solid, in methylene chloride is washed with water and dilute sodium chloride, dried (Na_2SO_4) and concentrated to give 7.25 g of the oxime, a compound of formula 6 (Scheme 1). A stirred solution of the oxime in acetone (165 mL), under nitrogen, is treated with 5% aqueous sodium carbonate (165 mL) and then dropwise during 20 minutes with a solution of *p*-toluenesulfonyl chloride (4.92 g, 0.0258 mol) in acetone (80 mL). The mixture is kept at ambient temperature for 18 hours and then concentrated under reduced pressure. A solution of the residue in methylene chloride is washed with water and dilute sodium chloride, dried (Na_2SO_4) and concentrated. Chromatography of the residue on silica gel with 3% methanol - 0.3% ammonium hydroxide-methylene chloride gave 5.98 g of the titled compound.

Step 2: Preparation of (*S*)-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]amine.

An ice cold, stirred mixture of the product of Example 2, Step 1 (0.22 g, 0.50 mmol) in tetrahydrofuran (THF; 15 mL), under nitrogen, is treated dropwise during 2 minutes, with a 1M solution of tetrabutylammonium fluoride in THF (1.5 mL). The mixture is kept in the ice bath for 10 minutes and at ambient temperature (24 °C) for 1 hour 25 minutes, diluted with ethyl acetate, washed with water and brine, dried (Na_2SO_4) and concentrated. Chromatography of the residue on silica gel with mixtures of methanol-methylene chloride containing 3-6% methanol gave 0.15 g of the alcohol, a compound of formula 9 (Scheme 1) where R is hydrogen: MS(ES) *m/z* 324 ($\text{M}+\text{H}^+$). A stirred suspension of the alcohol (0.15 g, 0.46 mmol) in methylene chloride (15 mL) and THF (8 mL), under nitrogen, is treated with triethylamine (0.5 mL, 1.4 mmol) and then portionwise during 1 minute at ambient temperature, with 0.14 g (0.56 mmol) of *m*-nitrobenzenesulfonyl chloride. The mixture is stirred for 90 minutes, mixed with additional methylene chloride (10 mL) to give a solution and kept at ambient temperature for 1 hour. It is then kept for several days at -11 °C, diluted with methylene chloride, washed with saturated sodium bicarbonate, water and brine, dried (Na_2SO_4) and


concentrated to give 0.21 g of the *m*-nitrobenzenesulfonate, a compound of formula 10 (Scheme 1). A stirred mixture of the *m*-nitrobenzenesulfonate (0.21 g, 0.44 mmol), acetonitrile (10 mL), 2-propanol (10 mL) and 29% ammonium hydroxide (10 mL) is warmed at 45-50 °C under a Dry Ice-acetone condenser for 4.5 hours and kept at ambient temperature for 18 hours. Additional ammonium hydroxide (5 mL) is added and the mixture is warmed at 45-50 °C for 4.5 hours, kept at ambient temperature for 1 hour, treated with 5 mL of ammonium hydroxide and kept at ambient temperature for 18 hours. It is then concentrated to give a yellow solid which is chromatographed on silica gel with mixtures of methanol-methylene chloride containing 5-7.5% methanol followed by 8% methanol-0.2% ammonium hydroxide-methylene chloride to give the titled product.

Step 3: Preparation of (*S*)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]thioacetamide.

A stirred solution of 0.12 g of the product of Example 2, Step 2 and 0.40 mL of triethylamine in a mixture of methylene chloride (10 mL) and methanol (10 mL), under nitrogen, is treated with ethyl dithioacetate (0.05 mL) and kept at ambient temperature for 145 hours. Additional 0.05 mL portions of ethyl dithioacetate are added after 24, 31 and 49 hours; additional triethylamine (1.0 mL) is also added after 49 hours. The mixture is concentrated to a small volume, diluted with ethyl acetate, washed with water and brine, dried (Na₂SO₄) and concentrated. Chromatography of the residue on silica gel with 3.5% methanol-methylene chloride gave 0.061 g of the titled product.

¹H NMR [300 MHz, (CD₃)₂SO] δ 2.42 (s, 3H), 2.56 (m, 2H), 3.07 (m, 4H), 3.24 (m, 2H), 3.76 (dd, 1H), 3.87 (m, 2H), 4.11 (t, 1H), 4.91 (m, 1H), 7.12 (m, 2H), 7.46 (dd, 1H), 7.67 (broad s, 1H), 10.35 (broad s, 1H).

EXAMPLE 3: Preparation of (*S*)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-4-methyl-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]-acetamide.



Step 1: Preparation of (*S*)-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-4-methyl-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]amine.

A mixture of 0.63 g (1.4 mmol) of the product of Example 2, Step 2, methyl iodide (0.093 mL) and THF (40 mL) is added dropwise during 12 minutes, under
5 nitrogen, to a stirred mixture of powdered potassium hydroxide (0.12 g) and tetrabutylammonium bromide (0.096 g) in THF (10 mL) and kept at ambient temperature for 20 hours. It is then diluted with ethyl acetate, washed with water and brine, dried (MgSO₄) and concentrated. Chromatography of the residue on silica gel with mixtures of acetone-methylene chloride that contained 10-40% acetone gave 0.46 g (71%) of the
10 methylated product, a compound of formula 8 (Scheme 1) where R' is methyl. An ice cold, stirred mixture of this product (0.17 g, 0.38 mmol) and THF (12 mL), under nitrogen, is treated dropwise with a 1M solution of tetrabutylammonium fluoride in THF (1.2 mL). It is kept in the ice bath for 15 minutes and at ambient temperature for 3 hours, mixed with ice water and extracted with ethyl acetate. The extract is washed with water
15 and brine, dried (MgSO₄) and concentrated to give 0.15 g of the alcohol, a compound of formula 9 (Scheme 1). A stirred, ice cold solution of the alcohol (0.52 g, 1.5 mmol) and triethylamine (0.60 mL) in methylene chloride (45 mL) is treated portionwise during 5 minutes with *m*-nitrobenzenesulfonyl chloride (0.42 g). The mixture is kept in the ice bath for 15 minutes and at ambient temperature for 3 hours, diluted with methylene
20 chloride, washed with saturated sodium bicarbonate, water and brine, dried (MgSO₄) and concentrated to give the *m*-nitrobenzenesulfonate, a compound of formula 10 (Scheme 1). A stirred mixture of this product, acetonitrile (35 mL), 2-propanol (35 mL) and concentrated ammonium hydroxide (35 mL) is kept at 45-50 °C under a Dry Ice-acetone condenser for 4.5 hours and at ambient temperature for 20 hours. Additional ammonium
25 hydroxide (6 mL) is added and the mixture is kept at 45-50 °C for 5.5 hours and at ambient temperature for 18 hours. The mixture is then concentrated under reduced pressure to remove the organic solvents and the aqueous residue is extracted first with ethyl acetate and then methylene chloride. The extracts are washed with water and brine, dried (MgSO₄) and concentrated. Chromatography of the residue on silica gel with
30 mixtures of methanol-methylene chloride containing 7.5-10% methanol gave the titled compound.

Step 2: Preparation of (*S*)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-4-methyl-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide.


A stirred, ice cold mixture of 0.10 g (0.30 mmol) of the product of Example 3, Step 1, and pyridine (1.74 mL), under nitrogen, is treated dropwise with acetic anhydride (0.57 mL, 6.04 mmol) and kept in the ice bath for 15 minutes and at ambient temperature for 3.5 hours. It is then concentrated in vacuo; the residue is mixed with ice water and saturated sodium bicarbonate and extracted with ethyl acetate. The extract is washed with water and brine, dried (MgSO₄) and concentrated. Crystallization of the residue from ethyl acetate-methanol gave 0.053 g of the titled compound.

mp 203-204 °C.

MS(ES) *m/z* 379 (M+H⁺), 401 (M+Na⁺).

Anal. calcd for C₁₈H₂₃FN₄O₄: C, 57.13; H, 6.13; N, 14.81. Found: C, 57.05; H, 6.23; N, 14.85.

EXAMPLE 4 Preparation of (*S*)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-4-methyl-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]-thioacetamide.



An ice cold, stirred solution of 0.18 g (0.535 mmol) of the product of Example 3, Step 1 and triethylamine (0.21 mL) in THF (8 mL) and methylene chloride (10 mL) is treated with a solution of ethyl dithioacetate (0.074 mL, 0.64 mmol) in THF (2 mL). The mixture is kept at ambient temperature for 20 hours, treated with one drop of additional ethyl dithioacetate and kept at ambient temperature for 7 hours. It is then concentrated under a stream of nitrogen. The residue is mixed with methylene chloride, washed with saturated sodium bicarbonate, water and brine, dried (Na₂SO₄) and concentrated. Chromatography of the residue on silica gel with mixtures of methanol-methylene chloride containing 2-4% methanol and crystallization of the product from ethyl acetate gave 0.13 g of the titled compound. mp 157-158 °C.

Anal. calcd for C₁₈H₂₃FN₄O₃S: C, 54.81; H, 5.88; N, 14.20. Found: C, 54.83; H, 5.93; N, 14.11.

EXAMPLE 5 MIC Test Method

The *in vitro* MICs of test compounds are determined by a standard agar dilution method. A stock drug solution of each analog is prepared in the preferred solvent, usually DMSO:H₂O (1:3). Serial 2-fold dilutions of each sample are made using 1.0 ml aliquots of sterile distilled water. To each 1.0 ml aliquot of drug is added 9 ml of molten Mueller Hinton agar medium. The drug-supplemented agar is mixed, poured into 15 x 100 mm petri dishes, and allowed to solidify and dry prior to inoculation.

Vials of each of the test organisms are maintained frozen in the vapor phase of a liquid nitrogen freezer. Test cultures are grown overnight at 35 C on the medium appropriate for the organism. Colonies are harvested with a sterile swab, and cell suspensions are prepared in Trypticase Soy broth (TSB) to equal the turbidity of a 0.5 McFarland standard. A 1:20 dilution of each suspension is made in TSB. The plates containing the drug supplemented agar are inoculated with a 0.001 ml drop of the cell suspension using a Steers replicator, yielding approximately 10⁴ to 10⁵ cells per spot. The plates are incubated overnight at 35 C.

Following incubation the Minimum Inhibitory Concentration (MIC µg/ml), the lowest concentration of drug that inhibits visible growth of the organism, is read and recorded. The data is shown in Table I.

TABLE I

Example No.	SAUR ^a 9213 MIC	SEPI ^b 12084 MIC	EFAE ^c 9217 MIC	SPNE ^d 9912 MIC	HINF ^e 30063 MIC	MCAT ^f 30610 MIC
1	4	1	4	0.5	8	8
3	4	1	4	1	32	8
4	1	<0.5	1	<0.5	8	2


20

- a) *S. aureus*, culture 9213
- b) *S. epidermidis*, culture 12084
- c) *E. faecalis*, culture 9217
- d) *S. pneumoniae*, culture 9912
- e) *H. influenzae*, culture 30063
- f) *M. catarrhalis*, culture 30610

25

CLAIMS

1. A compound of Formula I:



5

I

or a pharmaceutically acceptable salt thereof wherein:

R is H, C₂₋₆ alkenyl, C₂₋₇ alkynyl, C₁₋₆ alkyl, or C₁₋₆ alkyl substituted with one or two of the following:

- 10 a) F,
 b) Cl,
 c) CF₃,
 d) -OH,
 e) C₁₋₄ alkoxy,
 15 f) -CH₂C(=O)C₁₋₄ alkyl,
 g) -OC(=O)N(R₄)₂,
 h) C₁₋₄ alkyl S(O)_n, (wherein n is 0, 1 or 2),
 i) -CN,
 j) carboxy,
 20 k) -C₁₋₄ alkoxycarbonyl,
 l) -C(=O)N(R₄)₂,
 m) -N(R₄)SO₂C₁₋₄ alkyl,
 n) -N(R₄)C(=O)C₁₋₄ alkyl,
 o) -N(R₄)C(=O)N(R₄)₂,
 25 p) -N(R₄)C(=O)C₁₋₄ alkoxy,
 q) aryl, or
 r) Het;

aryl is phenyl, optionally substituted with one or two of the following:

- 30 a) F,
 b) Cl,

- c) Br,
 d) -CF₃,
 e) CN,
 f) C₁₋₃ alkoxy, or
 5 g) C₁₋₃ alkylthio;

Het is a 5- or 6-membered heteroaromatic moiety having 1-3 N, O or S atoms, optionally substituted with the following:

- a) F,
 b) Cl,
 10 c) C₁₋₃ alkoxy,
 d) C₁₋₃ alkylthio, or
 e) CN;


R₁ and R₂ are independently

- a) H,
 15 b) F, or
 c) Cl;

R₃ is

- a) C₁₋₆ alkyl, optionally substituted with one to three F or one to two Cl,
 b) C₁₋₆ alkoxy,
 20 c) amino,
 d) C₁₋₆ alkylamino,
 e) C₁₋₆ dialkylamino
 f) C₃₋₆ cycloalkyl,
 g) C₁₋₆ alkylthio, or

25

- h)  (wherein m is 0, 1, 2, 3 or 4);

R₄ is

- a) H, or
 b) C₁₋₃ alkyl; and
 30 X is O or S.

2. A compound of Claim 1 wherein X is O.

3. A compound of Claim 1 wherein X is S.
4. A compound of Claim 1 wherein R is H.
5. A compound of Claim 1 wherein R is C₁₋₄ alkyl.
6. A compound of Claim 1 wherein R₃ is C₁₋₄ alkyl, optionally substituted with one to three F or one to two Cl.
7. The compound of Claim 1 wherein Formula I is the S-enantiomer.
8. A compound of Claim 1 which is
- (a) (S)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide,
- (b) (S)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]thioacetamide,
- (c) (S)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-4-methyl-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]acetamide, or
- (d) (S)-N-[[3-[3-fluoro-4-(1,2,3,4,6,7-hexahydro-4-methyl-5-oxo-1,4-diazepin-1-yl)phenyl]-2-oxo-5-oxazolidinyl]methyl]thioacetamide.
9. A method for treating microbial infections in human comprising administering to a patient in need thereof an effective amount of a compound of Formula I as shown in Claim 1.
10. A pharmaceutical composition comprising a compound of Formula I as shown in Claim 1 and a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/22639

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D413/10 A61K31/42 C07D413/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 27188 A (ZENECA LTD) 31 July 1997 cited in the application see claims ---	1-10
Y	WO 95 25106 A (UPJOHN CO) 21 September 1995 see claims ---	1-10
Y	WO 93 23384 A (UPJOHN CO) 25 November 1993 cited in the application see claims ---	1-10
Y	WO 95 14684 A (UPJOHN CO) 1 June 1995 see claims -----	1-10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

4 February 1999

Date of mailing of the international search report

12/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/22639

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9727188 A	31-07-1997	AU 1315797 A EP 0882041 A	20-08-1997 09-12-1998
WO 9525106 A	21-09-1995	JP 8073455 A AU 681953 B AU 2099995 A CA 2183972 A CN 1143961 A EP 0750618 A JP 9512785 T NZ 283011 A US 5668286 A	19-03-1996 11-09-1997 03-10-1995 21-09-1995 26-02-1997 02-01-1996 22-12-1997 22-09-1997 16-09-1997
WO 9323384 A	25-11-1993	AU 668733 B AU 4287793 A CA 2133079 A CN 1079964 A CZ 9402505 A EP 0640077 A FI 945246 A HU 72296 A HU 9500659 A IL 105555 A JP 7506829 T MX 9302665 A NO 944237 A PL 174909 B PL 174850 B SK 133794 A US 5547950 A US 5700799 A ZA 9302855 A	16-05-1996 13-12-1993 25-11-1993 29-12-1993 16-08-1995 01-03-1995 08-11-1994 29-04-1996 28-11-1995 15-07-1998 27-07-1995 01-11-1993 04-01-1995 30-10-1998 30-09-1998 07-06-1995 20-08-1996 23-12-1997 24-10-1994
WO 9514684 A	01-06-1995	AU 698699 B AU 8010394 A CA 2174107 A CN 1135752 A EP 0730591 A JP 9505582 T NZ 274966 A US 5652238 A ZA 9407885 A	05-11-1998 13-06-1995 01-06-1995 13-11-1996 11-09-1996 03-06-1997 26-01-1998 29-07-1997 09-04-1996

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/22639

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 9
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 9 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.